Brain Research, 556 (1991) 358–359
Elsevier

Author Index

Adler, M., see Sheridan, R.E., 53
Aghajanian, G.K., see Alreja, M., 339
Akaike, N., Takahashi, K. and Morimoto, M., Heterogeneous distribution of tetrodotoxin-sensitive calcium-conducting channels in rat hippocampal CA1 neurons, 135
Alberch, J., see Arenas, E., 349
Alcayaga, J., see Gual, A., 139
Alreja, M. and Aghajanian, G.K., Pacemaker activity of locus coeruleus neurons: whole-cell recordings in brain slices show dependence on cAMP and protein kinase A, 339
Amir, S., De Blasio, E. and English, A.M., N^G-Monomethyl-L-arginine co-injection attenuates the thermogenic and hyperthermic effects of E_2 prostaglandin microinjection into the anterior hypothalamic preoptic area in rats, 157
Angelaki, D.E., see Dickman, J.D., 303
Arenas, E., Alberch, J., Pérez-Navarro, E., Solsona, C. and Marsal, J., Neostriatal dopaminergic terminals prevent the GABAergic involvement in the μ- and δ-opioid inhibition of KCl-evoked endogenous acetylcholine release, 349
Asai, K., Hotta, T., Nakanishi, K., Ito, J.-i., Tanaka, R., Otsuka, T., Matsui, N. and Kato, T., von Recklinghausen neurofibroma produces neuronal and glial growth-modulating factors, 344
Asanuma, M., see Ogawa, N., 271
Ault, B., see Caviedes, P., 285

Balin, B.J. and Lee, V.M-Y., Individual neurofilament subunits reassembled in vitro exhibit unique biochemical, morphological and immunological properties, 196
Balin, B.J., Clark, E.A., Trojanowski, J.Q. and Lee, V.M-Y., Neurofilament reassembly in vitro: biochemical, morphological and immuno-electron microscopic studies employing monoclonal antibodies to defined epitopes, 181
Ben-Jonathan, N., see Garris, P.A., 123
Berretta, S., Perciavalle, V. and Poppele, R.E., Origin of cuneate projections to the anterior and posterior lobes of the rat cerebellum, 297
Björklund, A., see Fischer, W., 44
Bledsoe, S., see Zhou, F.C., 108
Bloedel, J.R., see Bracha, V., 317
Bracha, V., Wu, J.-Z., Cartwright, S. and Bloedel, J.R., Selective involvement of the spinal trigeminal nucleus in the conditioned nictitating membrane reflex of the rabbit, 317
Brandão, M.L., Lopez-Garcia, J.A., Graeff, F.G. and Roberts, M.H.T., Electrophysiological evidence for excitatory 5-HT_2 and depressant 5-HT_{1A} receptors on neurones of the rat midbrain tectum, 259

Calverley, R.K.S., see Jones, D.G., 247
Carlen, P.L., see Jahromi, S.S., 33
Cartwright, S., see Bracha, V., 317
Caviedes, P., Koistinaho, J., Ault, B. and Rapoport, S.I., Effects of nerve growth factor on electrical membrane properties of cultured dorsal root ganglia neurons from normal and trisomy 21 human fetuses, 285
Chan, R.K.W., Chan, Y.S. and Wong, T.M., Responses of cardiovascular neurons in the rostral ventrolateral medulla of the normotensive Wistar Kyoto and spontaneously hypertensive rats to iontophoretic application of angiotensin II, 145
Chan, Y.S., see Chan, R.K.W., 145
Chen, Z., Engberg, G., Hedner, T. and Hedner, J., Antagonistic effects of somatostatin and substance P on respiratory regulation in the rat ventrolateral medulla oblongata, 13
Chesler, M., Sakatani, K. and Hassan, A.Z., Elevation and clearance of extracellular K^+ following contusion of the rat spinal cord, 71
Clark, E.A., see Balin, B.J., 181
Coirini, H., see McCarthy, M.M., 321
Correia, M.J., see Dickman, J.D., 303
Cowell, J., see Luine, V., 171
Coyle, J.T., see Tsai, G., 151
Cruccu, G., Ferracuti, S., Leardi, M.G., Fabbri, A. and Manfredi, M., Nociceptive quality of the orbicularis oculi reflexes as evaluated by distinct opiate- and benzodiazepine-induced changes in man, 209

Dacko, S., see Schneider, J.S., 292
De Groat, W.C., see Kruse, M.N., 6
De Vlieger, T.A., see Wildering, W.C., 324
De Blasio, E., see Amir, S., 157
Deshpande, S.S., see Sheridan, R.E., 53
Dickman, J.D., Angelaki, D.E. and Correia, M.J., Response properties of gerbil otolith afferents to small angle pitch and roll tilts, 303
Dun, N.J., Wu, S.Y. and Lin, C.W., Excitatory effects of cholecystokinin octapeptide on rat nodose ganglion cells in vitro, 161

Engberg, G., see Chen, Z., 13
English, A.M., see Amir, S., 157
Eugenín, J., see Gual, A., 139
Evans, C., see Racine, R.J., 218
Eyzaguirre, C., see Gual, A., 139
Eyzaguirre, C., see He, S.-F., 333

Fabbri, A., see Cruccu, G., 209
Ferracuti, S., see Cruccu, G., 209
Fischer, W., Nilsson, O.G. and Björklund, A., In vivo acetylcholine release as measured by microdialysis is unaltered in the hippocampus of cognitively impaired aged rats with degenerative changes in the basal forebrain, 44
Fisher, M., see Wong, V.L.Y., 1
Frankfurt, M., see Luine, V., 171

Garris, P.A. and Ben-Jonathan, N., Effects of reuptake inhibitors on dopamine release from the stalk-median eminence and posterior pituitary in vitro, 123
Gebber, G.L., see Zhong, S., 130
Gerfen, C.R., Substance P (neurokinin-1) receptor mRNA is selectively expressed in cholinergic neurons in the striatum and basal forebrain, 165
Graeff, F.G., see Brandão, M.L., 259
Gual, A., Eugenín, J., Alcayaga, J., Stensaas, L.J. and Eyzaguirre, C., The chick chorioallantoic membrane promotes survival of co-transplanted rat carotid bodies and nodose ganglia, 139
Guyenet, P.G., see Sun, M.-K., 61

Haba, K., see Ogawa, N., 271
Hammang, J.P. and Messing, A., Immortalized retinal neurons used as immunogen for the generation of cell-specific antisera, 85
Hasegawa, M., see Houdou, S., 267
Hashimoto, T., see Houdou, S., 267
Hassan, A.Z., see Chesler, M., 71
Haykal-Coates, N., O'Callaghan, J.P., Reinhard, Jr., J.F. and Jensen, K.F., Pargyline and γ-butyrolactone enhance tyrosine hydroxylase immunostaining of nigrostriatal axons, 353
He, S.-F., Wei, J.-Y. and Eyzaguirre, C., Effects of relative hypoxia and hypercapnia on intracellular pH and membrane potential of cultured carotid body glomus cells, 333
Hedner, J., see Chen, Z., 13
Hedner, T., see Chen, Z., 13
Hedreen, J.C., see Tsai, G., 151
Hoebel, B.G., see West, H.L., 95
Hofman, F.M., see Wong, V.L.Y., 1
Hotta, T., see Asai, K., 344
Houdou, S., Kuruta, H., Hasegawa, M., Konomi, H., Takashima, S., Suzuki, Y. and Hashimoto, T., Developmental immunohistochemistry of catalase in the human brain, 267

Ishii, H., see Wong, V.L.Y., 1
Ito, J.-i., see Asai, K., 344

Jahromi, S.S., Niesen, C. and Carlen, P.L., Amphetamine actions on pre- and postpubertal rat hippocampal dentate gran-

Brain Research, 556 (1991) 358–359
Elsevier

Author Index

Adler, M., see Sheridan, R.E., 53
Aghajanian, G.K., see Alreja, M., 339
Akaike, N., Takahashi, K. and Morimoto, M., Heterogeneous distribution of tetrodotoxin-sensitive calcium-conducting channels in rat hippocampal CA1 neurons, 135
Alberch, J., see Arenas, E., 349
Alcayaga, J., see Gual, A., 139
Alreja, M. and Aghajanian, G.K., Pacemaker activity of locus coeruleus neurons: whole-cell recordings in brain slices show dependence on cAMP and protein kinase A, 339
Amir, S., De Blasio, E. and English, A.M., N^G-Monomethyl-L-arginine co-injection attenuates the thermogenic and hyperthermic effects of E_2 prostaglandin microinjection into the anterior hypothalamic preoptic area in rats, 157
Angelaki, D.E., see Dickman, J.D., 303
Arenas, E., Alberch, J., Pérez-Navarro, E., Solsona, C. and Marsal, J., Neostriatal dopaminergic terminals prevent the GABAergic involvement in the μ- and δ-opioid inhibition of KCl-evoked endogenous acetylcholine release, 349
Asai, K., Hotta, T., Nakanishi, K., Ito, J.-i., Tanaka, R., Otsuka, T., Matsui, N. and Kato, T., von Recklinghausen neurofibroma produces neuronal and glial growth-modulating factors, 344
Asanuma, M., see Ogawa, N., 271
Ault, B., see Caviedes, P., 285

Balin, B.J. and Lee, V.M-Y., Individual neurofilament subunits reassembled in vitro exhibit unique biochemical, morphological and immunological properties, 196
Balin, B.J., Clark, E.A., Trojanowski, J.Q. and Lee, V.M-Y., Neurofilament reassembly in vitro: biochemical, morphological and immuno-electron microscopic studies employing monoclonal antibodies to defined epitopes, 181
Ben-Jonathan, N., see Garris, P.A., 123
Berretta, S., Perciavalle, V. and Poppele, R.E., Origin of cuneate projections to the anterior and posterior lobes of the rat cerebellum, 297
Björklund, A., see Fischer, W., 44
Bledsoe, S., see Zhou, F.C., 108
Bloedel, J.R., see Bracha, V., 317
Bracha, V., Wu, J.-Z., Cartwright, S. and Bloedel, J.R., Selective involvement of the spinal trigeminal nucleus in the conditioned nictitating membrane reflex of the rabbit, 317
Brandão, M.L., Lopez-Garcia, J.A., Graeff, F.G. and Roberts, M.H.T., Electrophysiological evidence for excitatory 5-HT_2 and depressant 5-HT_{1A} receptors on neurones of the rat midbrain tectum, 259

Calverley, R.K.S., see Jones, D.G., 247
Carlen, P.L., see Jahromi, S.S., 33
Cartwright, S., see Bracha, V., 317
Caviedes, P., Koistinaho, J., Ault, B. and Rapoport, S.I., Effects of nerve growth factor on electrical membrane properties of cultured dorsal root ganglia neurons from normal and trisomy 21 human fetuses, 285
Chan, R.K.W., Chan, Y.S. and Wong, T.M., Responses of cardiovascular neurons in the rostral ventrolateral medulla of the normotensive Wistar Kyoto and spontaneously hypertensive rats to iontophoretic application of angiotensin II, 145
Chan, Y.S., see Chan, R.K.W., 145
Chen, Z., Engberg, G., Hedner, T. and Hedner, J., Antagonistic effects of somatostatin and substance P on respiratory regulation in the rat ventrolateral medulla oblongata, 13
Chesler, M., Sakatani, K. and Hassan, A.Z., Elevation and clearance of extracellular K^+ following contusion of the rat spinal cord, 71
Clark, E.A., see Balin, B.J., 181
Coirini, H., see McCarthy, M.M., 321
Correia, M.J., see Dickman, J.D., 303
Cowell, J., see Luine, V., 171
Coyle, J.T., see Tsai, G., 151
Cruccu, G., Ferracuti, S., Leardi, M.G., Fabbri, A. and Manfredi, M., Nociceptive quality of the orbicularis oculi reflexes as evaluated by distinct opiate- and benzodiazepine-induced changes in man, 209

Dacko, S., see Schneider, J.S., 292
De Groat, W.C., see Kruse, M.N., 6
De Vlieger, T.A., see Wildering, W.C., 324
De Blasio, E., see Amir, S., 157
Deshpande, S.S., see Sheridan, R.E., 53
Dickman, J.D., Angelaki, D.E. and Correia, M.J., Response properties of gerbil otolith afferents to small angle pitch and roll tilts, 303
Dun, N.J., Wu, S.Y. and Lin, C.W., Excitatory effects of cholecystokinin octapeptide on rat nodose ganglion cells in vitro, 161

Engberg, G., see Chen, Z., 13
English, A.M., see Amir, S., 157
Eugenín, J., see Gual, A., 139
Evans, C., see Racine, R.J., 218
Eyzaguirre, C., see Gual, A., 139
Eyzaguirre, C., see He, S.-F., 333

Fabbri, A., see Cruccu, G., 209
Ferracuti, S., see Cruccu, G., 209
Fischer, W., Nilsson, O.G. and Björklund, A., In vivo acetylcholine release as measured by microdialysis is unaltered in the hippocampus of cognitively impaired aged rats with degenerative changes in the basal forebrain, 44
Fisher, M., see Wong, V.L.Y., 1
Frankfurt, M., see Luine, V., 171

Garris, P.A. and Ben-Jonathan, N., Effects of reuptake inhibitors on dopamine release from the stalk-median eminence and posterior pituitary in vitro, 123
Gebber, G.L., see Zhong, S., 130
Gerfen, C.R., Substance P (neurokinin-1) receptor mRNA is selectively expressed in cholinergic neurons in the striatum and basal forebrain, 165
Graeff, F.G., see Brandão, M.L., 259
Gual, A., Eugenín, J., Alcayaga, J., Stensaas, L.J. and Eyzaguirre, C., The chick chorioallantoic membrane promotes survival of co-transplanted rat carotid bodies and nodose ganglia, 139
Guyenet, P.G., see Sun, M.-K., 61

Haba, K., see Ogawa, N., 271
Hammang, J.P. and Messing, A., Immortalized retinal neurons used as immunogen for the generation of cell-specific antisera, 85
Hasegawa, M., see Houdou, S., 267
Hashimoto, T., see Houdou, S., 267
Hassan, A.Z., see Chesler, M., 71
Haykal-Coates, N., O'Callaghan, J.P., Reinhard, Jr., J.F. and Jensen, K.F., Pargyline and γ-butyrolactone enhance tyrosine hydroxylase immunostaining of nigrostriatal axons, 353
He, S.-F., Wei, J.-Y. and Eyzaguirre, C., Effects of relative hypoxia and hypercapnia on intracellular pH and membrane potential of cultured carotid body glomus cells, 333
Hedner, J., see Chen, Z., 13
Hedner, T., see Chen, Z., 13
Hedreen, J.C., see Tsai, G., 151
Hoebel, B.G., see West, H.L., 95
Hofman, F.M., see Wong, V.L.Y., 1
Hotta, T., see Asai, K., 344
Houdou, S., Kuruta, H., Hasegawa, M., Konomi, H., Takashima, S., Suzuki, Y. and Hashimoto, T., Developmental immunohistochemistry of catalase in the human brain, 267

Ishii, H., see Wong, V.L.Y., 1
Ito, J.-i., see Asai, K., 344

Jahromi, S.S., Niesen, C. and Carlen, P.L., Amphetamine actions on pre- and postpubertal rat hippocampal dentate gran-

ule neurons, 33
Janse, C., see Wildering, W.C., 324
Jensen, K.F., see Haykal-Coates, N., 353
Jones, D.G. and Calverley, R.K.S., Perforated and non-perforated synapses in rat neocortex: three-dimensional reconstructions, 247

Kato, T., see Asai, K., 344
Kawata, M., see Ueda, S., 329
Kenney, M.J., see Zhong, S., 130
Koistinaho, J., see Caviedes, P., 285
Konomi, H., see Houdou, S., 267
Kruse, M.N., Mallory, B.S., Noto, H., Roppolo, J.R. and De Groat, W.C., Properties of the descending limb of the spinobulbospinal micturition reflex pathway in the cat, 6
Kuncl, R., see Tsai, G., 151
Kuruta, H., see Houdou, S., 267

Laihinen, A., see Rinne, J.O., 117
Lazdunski, M., see Moll, C., 311
Leardi, M.G., see Cruccu, G., 209
Lebeda, F.J., see Sheridan, R.E., 53
Lee, V.M-Y., see Balin, B.J., 181
Lee, V.M-Y., see Balin, B.J., 196
Lin, C.W., see Dun, N.J., 161
Lönnberg, P., see Rinne, J.O., 117
Lopez-Garcia, J.A., see Brandão, M.L., 259
Lovenberg, T.W., Nichols, D.E., Nestler, E.J., Roth, R.H. and Mailman, R.B., Guanine nucleotide binding proteins and the regulation of cyclic AMP synthesis in NS20Y neuroblastoma cells: role of D_1 dopamine and muscarinic receptors, 101
Luine, V., Cowell, J. and Frankfurt, M., GABAergic-serotonergic interactions in regulating lordosis, 171

Mailman, R.B., see Lovenberg, T.W., 101
Mallory, B.S., see Kruse, M.N., 6
Manfredi, M., see Cruccu, G., 209
Marjamäki, P., see Rinne, J.O., 117
Mark, G.P., see West, H.L., 95
Marsal, J., see Arenas, E., 349
Matsui, N., see Asai, K., 344
Matsumoto, R.R. and Walker, J.M., Inhibition of rubral neurons by noxious and non-noxious pressure, 78
McCarthy, M.M., Coirini, H., Schumacher, M., Pfaff, D.W., McEwen, B.S. and Schwartz-Giblin, S., Ovarian steroid modulation of [^{3}H]muscimol binding in the spinal cord of the rat, 321
McEwen, B.S., see McCarthy, M.M., 321
Messing, A., see Hammang, J.P., 85
Miguel-Hidalgo, J.J., Senba, E., Takatsuji, K. and Tohyama, M., Ultrastructure and retinal innervation of deafferentation-induced enkephalin-immunoreactive elements in the superficial layers of the rat superior colliculus, 175
Moll, C., Mourre, C., Lazdunski, M. and Ulrich, J., Increase of sodium channels in demyelinated lesions of multiple sclerosis, 311
Moore, K.-A., see Racine, R.J., 218
Moore, K.-A., see Racine, R.J., 226
Mori, A., see Ogawa, N., 271
Morimoto, M., see Akaike, N., 135
Mourre, C., see Moll, C., 311
Murphy, J., see Zhou, F.C., 108

Nakanishi, K., see Asai, K., 344
Nestler, E.J., see Lovenberg, T.W., 101
Nichols, D.E., see Lovenberg, T.W., 101
Niesen, C., see Jahromi, S.S., 33
Nilsson, O.G., see Fischer, W., 44
Noto, H., see Kruse, M.N., 6

O'Callaghan, J.P., see Haykal-Coates, N., 353
Ogawa, N., Haba, K., Asanuma, M. and Mori, A., Long-lasting effect of cerulетide on dyskinesia and monoaminergic neuronal pathways in rats treated with iminodipropionitrile, 271
Otsuka, T., see Asai, K., 344

Perciavalle, V., see Berretta, S., 297
Pérez-Navarro, E., see Arenas, E., 349
Pfaff, D.W., see McCarthy, M.M., 321
Poppele, R.E., see Berretta, S., 297

Racine, R.J., Moore, K.-A. and Evans, C., Kindling-induced potentiation in the piriform cortex, 218
Racine, R.J., Moore, K.-A. and Wicks, S., Activation of the NMDA receptor: a correlate in the dentate gyrus field potential and its relationship to long-term potentiation and kindling, 226
Rapoport, S.I., see Caviedes, P., 285
Ray, P.E., Ruley, E.J. and Saavedra, J.M., Different effects of chronic K^+ depletion on forebrain and peripheral angiotensin II receptors in young rats, 240
Reinhard, Jr., J.F., see Haykal-Coates, N., 353
Rinne, J.O., Laihinen, A., Lönnberg, P., Marjamäki, P. and Rinne, U.K., A post-mortem study on striatal dopamine receptors in Parkinson's disease, 117
Rinne, U.K., see Rinne, J.O., 117
Roberts, M.H.T., see Brandão, M.L., 259
Romero, A., see Steinacker, A., 22
Roppolo, J.R., see Kruse, M.N., 6
Roth, R.H., see Lovenberg, T.W., 101
Rothstein, J.D., see Tsai, G., 151
Ruley, E.J., see Ray, P.E., 240

Saavedra, J.M., see Ray, P.E., 240
Sakatani, K., see Chesler, M., 71
Sano, Y., see Ueda, S., 329
Schneider, J.S. and Dacko, S., Relative sparing of the dopaminergic innervation of the globus pallidus in monkeys made hemi-parkinsonian by intracarotid MPTP infusion, 292
Schumacher, M., see McCarthy, M.M., 321
Schwartz-Giblin, S., see McCarthy, M.M., 321
Senba, E., see Miguel-Hidalgo, J.J., 175
Sheridan, R.E., Deshpande, S.S., Lebeda, F.J. and Adler, M., The effects of pumiliotoxin-B on sodium currents in guinea pig hippocampal neurons, 53
Sim, L., see Tsai, G., 151
Sircar, R. and Zukin, S.R., Kinetic mechanisms of glycine requirement for *N*-methyl-D-aspartate channel activation, 280
Solsona, C., see Arenas, E., 349
Stauch-Slusher, B., see Tsai, G., 151
Steinacker, A. and Romero, A., Characterization of voltage-gated and calcium-activated potassium currents in toadfish saccular hair cells, 22
Stensaas, L.J., see Gual, A., 139
Stornetta, R.L., see Sun, M.-K., 61
Sun, M.-K., Stornetta, R.L. and Guyenet, P.G., Morphology of rostral medullary neurons with intrinsic pacemaker activity in the rat, 61
Suzuki, Y., see Houdou, S., 267

Takahashi, K., see Akaike, N., 135
Takashima, S., see Houdou, S., 267
Takatsuji, K., see Miguel-Hidalgo, J.J., 175
Tanaka, R., see Asai, K., 344
Tohyama, M., see Miguel-Hidalgo, J.J., 175
Trojanowski, J.Q., see Balin, B.J., 181
Tsai, G., Stauch-Slusher, B., Sim, L., Hedreen, J.C., Rothstein, J.D., Kuncl, R. and Coyle, J.T., Reductions in acidic amino acids and *N*-acetylaspartylglutamate in amyotrophic lateral sclerosis CNS, 151

Ueda, S., Sano, Y. and Kawata, M., Collateral sprouting of serotonergic fibers in the cingulate cortex and the septum following cortical-hippocampal lesions, 329
Ulrich, J., see Moll, C., 311

Walker, J.M., see Matsumoto, R.R., 78
Wei, J.-Y., see He, S.-F., 333
West, H.L., Mark, G.P. and Hoebel, B.G., Effects of conditioned taste aversion on extracellular serotonin in the lateral hypothalamus and hippocampus of freely moving rats, 95
Wicks, S., see Racine, R.J., 226
Wildering, W.C., Janse, C. and De Vlieger, T.A., The role of pacemaker properties and synaptic input in generation and modulation of spiking activity in a pair of electrically coupled peptidergic neurons, 324
Wong, T.M., see Chan, R.K.W., 145
Wong, V.L.Y., Hofman, F.M., Ishii, H. and Fisher, M., Regional distribution of thrombomodulin in human brain, 1
Wu, J.-Z., see Bracha, V., 317
Wu, S.Y., see Dun, N.J., 161

Zhong, S., Kenney, M.J. and Gebber, G.L., High power, low frequency components of cardiac, renal, splenic and vertebral sympathetic nerve activities are uniformly reduced by spinal cord transection, 130
Zhou, F.C., Bledsoe, S. and Murphy, J., Serotonergic sprouting is induced by dopamine-lesion in substantia nigra of adult rat brain, 108
Zukin, S.R., see Sircar, R., 280